

SCIP Prophylactic Antibiotics (Regimen for Selected Surgeries)

<p><u>CABG, Other Cardiac Surgery, or Vascular Surgery</u></p> <p>Cefazolin, Cefuroxime, or Vancomycin**</p> <p>If beta-lactam allergy: Vancomycin* or Clindamycin*</p>	<p><u>Hip/Knee Arthroplasty</u></p> <p>Cefazolin, Cefuroxime, or Vancomycin**</p> <p>If beta-lactam allergy: Vancomycin* or Clindamycin*</p>
<p><u>Colon Surgery</u></p> <p>Cefotetan, Cefoxitin, or Ampicillin/Sulbactam</p> <p>Or</p> <p>Cefazolin, or Cefuroxime, AND Metronidazole</p> <p>If beta-lactam allergy: Clindamycin AND Amikacin, Gentamicin, or Tobramycin</p> <p>Or</p> <p>Clindamycin AND Moxifloxacin or Ciprofloxacin</p> <p>Or</p> <p>Clindamycin AND Aztreonam</p> <p>Or</p> <p>Metronidazole AND Amikacin, Gentamicin, or Tobramycin</p> <p>Or</p> <p>Metronidazole AND Moxifloxacin or Ciprofloxacin</p>	<p><u>Hysterectomy</u></p> <p>Cefotetan, Cefazolin, Cefoxitin, Cefuroxime, or Ampicillin/Sulbactam</p> <p>If beta-lactam allergy: Clindamycin AND Amikacin, Gentamicin, or Tobramycin</p> <p>Or</p> <p>Clindamycin AND Moxifloxacin or Ciprofloxacin</p> <p>Or</p> <p>Clindamycin AND Aztreonam</p> <p>OR</p> <p>Metronidazole AND Amikacin, Gentamicin, or Tobramycin</p> <p>Or</p> <p>Metronidazole AND Moxifloxacin or Ciprofloxacin</p>

*For cardiac, orthopedic, and vascular surgery, if the patient is allergic to beta-lactam antibiotics, Vancomycin or Clindamycin are acceptable substitutes

**Vancomycin is acceptable with a Physician/APN/PA/Pharmacist documented justification for its use

Special Note: Utilize the specific SCIP surgery prophylactic antibiotic order forms located on the Meridian Intranet under Clinical Forms. These forms provide the most current recommended antibiotics and antibiotics available on formulary.